

Real forms of quantum orthogonal groups, q -Lorentz groups in any dimension

Paolo Aschieri

*Theoretical Physics Group, Physics Division
Lawrence Berkeley National Laboratory, 1 Cyclotron Road
Berkeley, California 94720, USA.*

Abstract

We review known real forms of the quantum orthogonal groups $SO_q(N)$. New $*$ -conjugations are then introduced and we construct all real forms of quantum orthogonal groups. We thus give an RTT formulation of the $*$ -conjugations on $SO_q(N)$ that is complementary to the $\mathcal{U}_q(g)$ $*$ -structure classification of Twietmeyer [5]. In particular we easily find and describe the real forms $SO_q(N-1,1)$ for any value of N . Quantum subspaces of the q -Minkowski space are analyzed.

MSC-class: 81R50; 17B37

e-mail: aschieri@lbl.gov

1 Introduction

In the last years there has been an intense study of noncommutative deformations of the Lorentz group. A deformed Lorentz group acts on a deformed noncommutative Minkowski spacetime that has thus the same symmetry richness as in the classical case. An important aspect is that non commuting coordinates give rise to indetermination relations and discretization, this might be a realistic picture of how spacetime behaves at short distances (see for example [1]).

There are many deformations of the Lorentz group. In this paper we are concerned with the standard FRT [2] deformation of the orthogonal groups. There, the complex orthogonal quantum groups $SO_q(N)$ are defined for any $N > 2$ and various real forms are studied, in particular $SO_q(2n, 1)$ with $q \in \mathbf{R}$. A new real form $SO_q(n+1, n-1)$ with $|q| = 1$ was then described in [3]. We present here the real forms $SO_q(2n-1, 1)$ with $q \in \mathbf{R}$ and therefore construct q -Lorentz groups in any dimensions. More in general, for $q \in \mathbf{R}$, we can obtain any desired signature, $SO_q(l, m)$, $l + m = N$ as well as deformations of the real form $SO^*(2n)$.

This letter is organized as follows. We first briefly recall the R -matrix construction of the orthogonal quantum groups, we then discuss in the undeformed case the equivalence between real forms and $*$ -structures (conjugations). It is the $*$ -structure formulation that has a straightforward generalization to the quantum case, see however [4] for a different approach. We end the section recalling some of the $*$ -conjugations defined in [2]. Section 3 contains the main results, we find an involution $\sharp$ that is an $SO_q(N)$ automorphism (algebra and coalgebra map compatible with the antipode of $SO_q(N)$). The composition $*^\sharp = \sharp \circ *$ of $\sharp$ with the $*$ -conjugations defined in [2] gives new star structures that lead to the real forms: $SO_q(2n-1, 1)$ with $q \in \mathbf{R}$, $SO_q(n, n+1)$ with $|q| = 1$ and $SO_q(n+1, n-1)$ with $|q| = 1$. The conjugations on the quantum orthogonal plane that are associated to these $*^\sharp$ -conjugations are also easily found. This procedure can be reiterated with another set of involutive automorphisms $\natural$, $\sharp \circ \natural$, that composed with $*$ give, for $q \in \mathbf{R}$, pseudo-orthogonal q -groups with any signature as well as the q -groups $SO_q^*(2n)$. Comparison of our results with [5] shows that via the involutive automorphisms $\sharp$, $\natural$ we obtain a complete classification of all the real forms of orthogonal q -groups. In Note 2 and Note 1 we remark that this is the same construction as in the classical case, where all real forms of $SO(N, \mathbf{C})$ are classified via involutive automorphism $\sharp$, $\natural$ of the compact real form $SO(N, \mathbf{R})$. These involutive automorphisms $\sharp$ ($\natural$) of $SO(N, \mathbf{R})$ can be realized via matrices $\mathcal{D} \in O(N, \mathbf{R})$: $\sharp : SO(N, \mathbf{R}) \rightarrow SO(N, \mathbf{R})$, $T^\sharp = \mathcal{D}T\mathcal{D}^{-1}$ with $\mathcal{D}^2 = \pm \mathbf{1}$. Similarly in the quantum case the $SO_q(N, \mathbf{R})$ involutive automorphisms $\sharp$, $\natural$ are realized via matrices (with complex entries) that satisfy the RTT and orthogonality relations of $SO_q(N)$, that have q -determinant $= \pm 1$, that squared equal $\pm \mathbf{1}$ and that moreover satisfy the reality conditions of $SO_q(N, \mathbf{R})$. In Note 2 we also explicitly classify the involutive automorphisms of $SO_q(N, \mathbf{R})$ ($\forall N \neq 8$).

We conclude by observing that even if the classical inclusion $SO(N) \subset SO(N+1)$ no more holds at the quantum level, one still has that the $(N+1)$ -dimensional q -

orthogonal plane includes the N -dimensional q -orthogonal plane. In particular the q -Minkowski spacetime contains the q -euclidean space.

2 $SO_q(N)$ and the real forms $SO_q(N, \mathbf{R})$, $SO_q(n, n)$, $SO_q(n+1, n)$

The quantum group $SO_q(N)$ is freely generated by the non-commuting matrix elements T^a_b (fundamental representation, $a, b = 1, \dots, N$) and the unit element I , modulo the quadratic RTT and CTT (orthogonality) relations discussed below and the condition $\det_q T = 1$ where $\det_q$ is the quantum determinant defined via the quantum epsilon tensor $\varepsilon_q^{i_1, \dots, i_N}$ in [6]. The noncommutativity is controlled by the R matrix: $R^{ab}_{ef} T^e_c T^f_d = T^b_f T^a_e R^{ef}_{cd}$ i.e.:

$$R_{12} T_1 T_2 = T_2 T_1 R_{12} . \quad (2.1)$$

The orthogonality relations are $C^{bc} T^a_b T^d_c = C^{ad} I$, $C_{ac} T^a_b T^c_d = C_{bd} I$ that we rewrite in matrix notation as:

$$TCT^t = CI \quad , \quad T^t CT = CI . \quad (2.2)$$

Following [2] we define prime indices as $a' \equiv N + 1 - a$. The indices run over N values, for N even we write $N = 2n$, otherwise $N = 2n + 1$, in this case we define $n_2 \equiv \frac{N+1}{2}$. Notice that for $N = 2n$ we have $n' = n + 1$. We also define a vector ρ_a as:

$$(\rho_1, \dots, \rho_N) = \begin{cases} (\frac{N}{2} - 1, \frac{N}{2} - 2, \dots, \frac{1}{2}, 0, -\frac{1}{2}, \dots, -\frac{N}{2} + 1) & \text{for } SO(2n+1) \\ (\frac{N}{2} - 1, \frac{N}{2} - 2, \dots, 1, 0, 0, -1, \dots, -\frac{N}{2} + 1) & \text{for } SO(2n) \end{cases} \quad (2.3)$$

Then the (antidiagonal) metric is :

$$C_{ab} = q^{-\rho_a} \delta_{ab'} \quad (2.4)$$

its inverse C^{ab} satisfies $C^{ab} C_{bc} = \delta_c^a = C_{cb} C^{ba}$, it is easily seen that the matrix elements of the metric and the inverse metric coincide.

The nonzero complex components of the R matrix are (no sum on repeated indices):

$$\begin{aligned} R^{aa}_{aa} &= q, & a &\neq n_2 \\ R^{aa'}_{aa'} &= q^{-1}, & a &\neq n_2 \\ R^{n_2 n_2}_{n_2 n_2} &= 1 \\ R^{ab}_{ab} &= 1, & a &\neq b, a' \neq b \\ R^{ab}_{ba} &= q - q^{-1}, & a &> b, a' \neq b \\ R^{aa'}_{a'a} &= (q - q^{-1})(1 - q^{\rho_a - \rho_{a'}}) = (q - q^{-1})[1 - C^{a'a} C_{a'a}], & a &> a' \\ R^{aa'}_{bb'} &= -(q - q^{-1})q^{\rho_a - \rho_b} = -(q - q^{-1})C^{a'a} C_{bb'}, & a &> b, a' \neq b \end{aligned} \quad (2.5)$$

The coproduct, counit and antipode are as usual defined by: $\Delta(T^a_b) = T^a_c \otimes T^c_b$, $\varepsilon(T^a_b) = \delta^a_b$ and $\kappa(T) = CT^tC$ i.e. $\kappa(T^a_b) = C^{ae}T^f_e C_{fb}$.

Quantum orthogonal planes are defined by the commutations

$$P_A^{ab}{}_{cd} x^c x^d = 0 \quad (2.6)$$

where the q -antisymmetrizer P_A is given by $P_A = \frac{1}{q+q^{-1}}[-\hat{R} + rI - (q - q^{1-N})P_0]$ and $P_0^{ab}{}_{cd} = (C_{ef}C^{ef})^{-1}C^{ab}C_{cd}$. The symmetry of the q -planes under the q -groups is expressed via the coaction δ : $\delta(x^a) = T^a_b \otimes x^b$.

Star Structures and Real Forms

The Hopf algebra $SO_q(N)$ can be interpreted as the deformation of the algebra of functions on a group manifold only introducing a $*$ -structure on $SO_q(N)$ (the analogue of complex conjugation). This procedure leads to the quantum groups $SO_q(N, \mathbf{R}) = Fun_q(SO(N, \mathbf{R}))$, $SO_q(N-1, 1) = Fun_q(SO(N-1, 1))$ and more in general $SO_q(l, m) = Fun_q(SO(l, m))$ and $SO_q^*(2n) = Fun_q(SO^*(2n))$. A $*$ -structure or $*$ -conjugation on a Hopf algebra A (and in particular on $SO_q(N)$) is an algebra anti-automorphism $(\eta ab)^* = \bar{\eta} b^* a^* \forall a, b \in A, \forall \eta \in \mathbf{C}$; coalgebra automorphism $\Delta \circ * = (* \otimes *) \circ \Delta$, $\varepsilon \circ * = \varepsilon$ and involution $*^2 = id$. It follows that $* \circ \kappa^{-1} = \kappa \circ *$ i.e. $[\kappa^{-1}(T)]^* = \kappa(T^*)$.¹ Two star structures $*$ and $'$ are equivalent if there exist a Hopf algebra automorphism α such that $' = \alpha \circ * \circ \alpha^{-1}$.

Let us clarify the relation between real forms of groups and $*$ -structures on Hopf algebras.

Consider a real form $g_{\mathbf{R}}$ of a complex Lie algebra g i.e. $g = g_{\mathbf{R}} \oplus i g_{\mathbf{R}}$. We exponentiate the elements of $g_{\mathbf{R}}$ to get the real form $G_{\mathbf{R}}$ of the complex Lie group G . There is a one-to-one correspondence between real forms of the complex Lie algebra g and star structures on the universal enveloping algebra $\mathcal{U}(g)$. The $*$ -operation that acts as minus the identity on $g_{\mathbf{R}}$ satisfies $[\chi, \chi']^* = [\chi'^*, \chi^*] \forall \chi, \chi' \in g_{\mathbf{R}}$ and is uniquely extended as an anti-linear, anti-multiplicative and involutive map on the Hopf algebra $\mathcal{U}(g)$. We also have a corresponding $*$ -structure on $\mathcal{F}(G)$, the algebra of analytic functions on the complex group G . We define $*$: $f \mapsto f^*$ such that $f^*(g) \equiv \overline{f(g)} \forall g \in G_{\mathbf{R}}$, then f^* is extended analytically to all G . Viceversa a $*$ -structure on $\mathcal{F}(G)$ determines the following real form $G_{\mathbf{R}}$ of G : $G_{\mathbf{R}} = \{g \in G \mid f^*(g) = \overline{f(g)}\}$. In other words, the algebra of regular functions on $G_{\mathbf{R}}$ [for ex. $SO(N, \mathbf{R})$] is isomorphic to the algebra of regular functions on $G_{\mathbf{R}'}$ [for ex. $SO(N-1, 1)$] and to the algebra of analytic functions on the complex manifold G [$SO(N, \mathbf{C})$]; indeed any regular function $f \in Fun(G_{\mathbf{R}})$ can be analytically continued in a unique function $\hat{f} \in \mathcal{F}(G)$, then the restriction of $\hat{f}$ to the $G_{\mathbf{R}'}$ sub-manifold of G belongs to $Fun(G_{\mathbf{R}'})$. Only considering a $*$ -structure we can understand if the orthogonal matrix entries T^a_b generate functions on $SO(N, \mathbf{R})$ or on $SO(N-1, 1)$.

¹Proof: $\delta^a_c I = \varepsilon(T^a_c) I = (\kappa \otimes id) \Delta(T^a_c) = \kappa(T^a_b) T^b_c$. Apply $*$ to get $\delta^a_c I = T^b_c [\kappa(T^a_b)]^*$; multiply from the left with $\kappa^{-1}(T^c_e)$ to get the thesis. $\square$

We have seen that a real form $g_{\mathbf{R}}$ of g is equivalent to a $*$ -structure on $\mathcal{U}(g)$ and to a $*$ -structure on $\mathcal{F}(G)$, the algebra of analytic functions on G . The $*$ -structures on $\mathcal{F}(G)$ and $\mathcal{U}(g)$ can be directly related using the duality $\langle \cdot, \cdot \rangle$ between these two Hopf algebras. Since $f^*(g) = \overline{f(g)} \forall g \in G_{\mathbf{R}}$ we have, write $g = e^{\chi}$, $\langle \chi, f^* \rangle = \overline{\langle \chi, f \rangle} \forall \chi \in g_{\mathbf{R}}$. More in general $\forall \psi \in \mathcal{U}(g), \forall f \in \mathcal{F}(G)$,

$$\langle \psi, f^* \rangle = \overline{\langle \kappa(\psi)^*, f \rangle}. \quad (2.7)$$

Proof: write

$$\langle \psi, f^* \rangle = \overline{\langle \mathcal{O}(\psi), f \rangle} \quad (2.8)$$

we want to find the map $\mathcal{O}$ implicitly defined by the above relation and such that $\mathcal{O}$ on $g_{\mathbf{R}}$ is the identity. From (2.8) it is easy to see that $\mathcal{O}$ is antilinear and multiplicative, this uniquely defines $\mathcal{O} = * \circ \kappa$, indeed $* \circ \kappa(\chi) = -\chi^* = \chi$. $\square$

Finally it is straightforward to see that if $*$ is equivalent to $*'$ i.e., if there exists an automorphism α of $\mathcal{U}(g)$ [or $\mathcal{F}(G)$] such that $*' = \alpha \circ * \circ \alpha^{-1}$, then $g_{\mathbf{R}}$ is isomorphic to $g_{\mathbf{R}'}$ and $G_{\mathbf{R}}$ is isomorphic to $G_{\mathbf{R}'}$; the viceversa holds as well.

In the quantum case, following [2], on orthogonal q -groups a conjugation can be defined

- trivially as $T^{\times} = T$ i.e. $T^a_b{}^{\times} = T^a_b$. Compatibility with the RTT relations (2.1) requires $\bar{R}_q = R_q^{-1} = R_{q^{-1}}$, i.e. $|q| = 1$. Then the CTT relations are invariant under $*$ -conjugation. The corresponding real forms are $SO_q(n, n; \mathbf{R})$, $SO_q(n+1, n; \mathbf{R})$. To prove this, recall that in the classical limit a $*$ structure on G determines the real form $G_{\mathbf{R}} = \{g \in G \mid f^*(g) = \overline{f(g)} \forall f \in \mathcal{F}(G)\}$; therefore in the $q \rightarrow 1$ limit the conjugation $*$ becomes the usual complex conjugation $\overline{}$ on functions on $G_{\mathbf{R}}$. In our case the condition $T^{\times}(g) = \overline{T(g)}$ reads $T(g) = \overline{T(g)}$. We have real matrices $T(g)$ orthogonal with respect to the metric $C_{q=1}^{ab} = \delta^{ab}$. In the new basis $T' = MT$, where $M^a_b = \frac{1}{\sqrt{2}}(\delta^a_b + \epsilon^a \delta^a_{b'})$ and $\epsilon^a = 1$ if $a < a'$, $\epsilon^a = -1$ if $a > a'$, $\epsilon^a = \sqrt{2} - 1$ if $a = a'$, we have $T'^t C' T' = C' I$ where $C' = (M^{-1})^t C M^{-1}$ is diagonal with signature (n, n) or $(n+1, n)$.

- via the metric as $T^* = [\kappa(T)]^t$ i.e. $T^* = C^t T C^t$. The condition on R is $\bar{R}^{ab}_{cd} = R^{dc}_{ba}$, which happens for $q \in \mathbf{R}$. Again the CTT relations are $\star$ -invariant. The metric on a real basis (we will see the explicit construction later on) has in the $q \rightarrow 1$ limit compact signature $(+, +, \dots)$ so that the real form is $SO_q(N; \mathbf{R})$.

The quantum orthogonal group co-acts on the quantum orthogonal plane, and may induce an associated conjugation on the q -plane as well. More precisely a conjugation on the quantum orthogonal plane —i.e. an anti-linear anti-multiplicative involution of the q -plane algebra— is compatible with a conjugation on its q -symmetry group if the coaction δ of the q -group on the q -plane: $\delta(x^a) = T^a_b \otimes x^b$ satisfies $\delta(x^*) = T^* \otimes x^* \equiv \delta^*(x)$. The above two q -groups conjugations have unique (up to an overall phase) associated q -plane conjugations, they are respectively $(x^a)^{\times} = x^a$ and $(x^a)^{\star} = C_{ba} x^b$.

3 Real forms $SO_q(N-1, 1)$, $SO_q(l, m)$ and $SO_q^*(2n)$

We first notice that if we have an involution $\sharp$ that is a Hopf algebra automorphism (algebra and coalgebra map compatible with the antipode: $\kappa(a^\sharp) = [\kappa(a)]^\sharp$) and that commutes with a conjugation $*$, then the composition of these two involutions: $*^\sharp \equiv \sharp \circ * = * \circ \sharp$ is again a conjugation. [Hint: it is trivially antilinear, antimultiplicative and compatible with the coproduct: $\Delta(T^{*\sharp}) = T^{*\sharp} \otimes T^{*\sharp}$. It is involutive because $(\sharp \circ *) \circ (\sharp \circ *) = id \Leftrightarrow \sharp \circ * = * \circ \sharp$]. We now find an involution $\sharp$ that comutes with $\times$ and $\star$ as defined above.

Define the map $\sharp$ on the generators of $SO_q(N)$ as:

$$T^\sharp = \mathcal{D} T \mathcal{D}^{-1} \quad \text{i.e.} \quad (T^a_b)^\sharp = \mathcal{D}^a_e T^e_f \mathcal{D}^f_b^{-1} \quad (3.9)$$

and extend it by linearity and multiplicativity to all $SO_q(N)$. The entries of the N -dimensional $\mathcal{D}$ matrix are

$$\mathcal{D} = \begin{pmatrix} 1 & & & & & \\ & \dots & & & & \\ & & 1 & & & \\ & & & 0 & 1 & \\ & & & 1 & 0 & \\ & & & & & 1 & \\ & & & & & & \dots & \\ & & & & & & & 1 \end{pmatrix}, \quad \mathcal{D} = \begin{pmatrix} 1 & & & & & \\ & \dots & & & & \\ & & 1 & & & \\ & & & -1 & & \\ & & & & 1 & \\ & & & & & \dots & \\ & & & & & & 1 \end{pmatrix} \quad (3.10)$$

for N even for N odd

In the $N = 2n$ case the $\mathcal{D}$ matrix exchanges the index n with the index $n+1$, in the $N = 2n+1$ case $\mathcal{D}$ change the sign of the entries of the T matrix as many times as the index $n_2 = \frac{N+1}{2}$ appears. Since $\mathcal{D}^2 = \mathbf{1}$ we immediately see that $\sharp$ is an involution.

We now prove that $\sharp$ is compatible with the algebra structure, i.e. it is compatible with the RTT and CTT relations. Use (2.5) to prove that

$$\mathcal{D}_1^{-1} \mathcal{D}_2^{-1} R_{12} \mathcal{D}_1 \mathcal{D}_2 = R_{12} \quad \text{i.e.} \quad R_{12} \mathcal{D}_1 \mathcal{D}_2 = \mathcal{D}_2 \mathcal{D}_1 R_{12}. \quad (3.11)$$

We also have

$$\mathcal{D}^t C \mathcal{D} = C \quad , \quad \mathcal{D} C \mathcal{D}^t = C \quad (3.12)$$

(the second equation in (3.12) follows from the first). Therefore $\mathcal{D}$ is a matrix of complex entries that satisfies the RTT and orthogonality relations. It is then trivial to show that $\sharp$ is compatible with the algebra structure, indeed $T^\sharp = \mathcal{D} T \mathcal{D}^{-1}$ is a product of q -orthogonal matrices whose matrix entries mutually commute, therefore $T^\sharp$ itself is a q -orthogonal matrix. We conclude that $R_{12} T_1^\sharp T_2^\sharp = T_2^\sharp T_1^\sharp R_{12}$, $T^\sharp C T^\sharp = C I$, $T^\sharp C T^\sharp = C I$ are equivalent to (2.1) and (2.2) and therefore $\sharp$ is an algebra automorphism of $SO_q(N)$. An explicit proof is also instructive, for example we have: $(T C T^t)^\sharp = \mathcal{D} T \mathcal{D} C \mathcal{D}^t T^t \mathcal{D}^t = \mathcal{D} (T C T^t) \mathcal{D}^t = \mathcal{D} C \mathcal{D}^t I = C I$.

The compatibility of $\sharp$ with the coalgebra structure is easily checked: $\Delta(T^\sharp) = \mathcal{D}\Delta(T)\mathcal{D}^{-1} = T^\sharp \otimes T^\sharp$, $\varepsilon(T^\sharp) = \varepsilon(T)$. Compatibility with the antipode: $\kappa(a^\sharp) = [\kappa(a)]^\sharp$ follows from compatibility with the algebra and coalgebra structure (cf. footnote 1). We now show that the two conjugations defined in the previous section commute with $\sharp$. For the second conjugation, defined by $T^\star = [\kappa(T)]^t = C^t T C^t$, we have $(T^\sharp)^\star = (T^\star)^\sharp \Leftrightarrow \overline{\mathcal{D}} C^t T C^t \overline{\mathcal{D}}^{-1} = C^t \mathcal{D} T \mathcal{D}^{-1} C^t$. For an arbitrary $\mathcal{D}$ matrix this last relation is equivalent to (use Schur lemma) $\overline{\mathcal{D}} C^t = \text{const} \cdot C^t \mathcal{D}$ i.e.

$$\overline{\mathcal{D}} = \text{const} \cdot C^t \mathcal{D} C^t . \quad (3.13)$$

In our case $\mathcal{D}$ is given by (3.10) and satisfies $\overline{\mathcal{D}} = C^t \mathcal{D} C^t$ so that $(T^\sharp)^\star = (T^\star)^\sharp$. The two maps $\star$ and $\sharp$ not only commute when applied to the T^a_b matrix entries, they also commute when applied to any element of the q -group because they are respectively multiplicative and antimultiplicative. For the first conjugation, defined by $T^\times = T$, the proof that $\sharp \circ \times = \times \circ \sharp$ is straightforward.

Associated to $\times^\sharp$ and $\star^\sharp$ we can construct the conjugations that act on the quantum orthogonal plane and are compatible with the coaction $x^a \rightarrow T^a_b \otimes x^b$. These conjugations respectively are: $(x^a)^{\times^\sharp} = \mathcal{D}^a_b x^b$ and $(x^a)^{\star^\sharp} = C_{ba} \mathcal{D}^b_e x^e$. Notice that these q -plane maps are well defined conjugations. This is so because $\sharp$, defined by $x^\sharp \equiv \mathcal{D}x$, is an involution: $\mathcal{D}^2 = \mathbf{1}$, and also on the q -plane $\sharp$ commutes with $\times$ or $\star$.

The above results are particular examples of the following theorem:

Theorem 1 Consider a map $\sharp$ on $SO_q(N)$ defined by (3.9) where $\mathcal{D}$ is an arbitrary matrix satisfying (3.11), (3.12) and $\mathcal{D}^2 = \pm \mathbf{1}$. Then $\sharp$ is an involutive automorphism of $SO_q(N)$. Moreover the compositions $\times^\sharp$ and $\star^\sharp$ of the conjugations $\times$ and $\star$ with the automorphism $\sharp$ is again a conjugation if and only if, respectively, $\sharp \circ \times = \times \circ \sharp$ and $\sharp \circ \star = \star \circ \sharp$. These commutations hold if and only if

$$\left. \begin{array}{l} \overline{\mathcal{D}} = \mathcal{D} \quad \text{with } \mathcal{D}^2 = \mathbf{1} \\ \overline{\mathcal{D}} = -\mathcal{D} \quad \text{with } \mathcal{D}^2 = -\mathbf{1} \end{array} \right\} \text{ for } \times , \quad (3.14)$$

$$\overline{\mathcal{D}} = C^t \mathcal{D} C^t \quad \text{for } \star . \quad (3.15)$$

If $\mathcal{D}^2 = \mathbf{1}$ then the associated maps $\times^\sharp$ or $\star^\sharp$ that act on the q -plane as:

$$x^{\star^\sharp} = C^t \mathcal{D} x \quad , \quad x^{\times^\sharp} = \mathcal{D} x \quad (3.16)$$

are well defined conjugations and are compatible with the q -group coaction.

Proof: The definition (3.9) and relations (3.11), (3.12) and $\mathcal{D}^2 = \pm \mathbf{1}$ assure that $\sharp$ is an automorphism and an involution of $SO_q(N)$. To prove that $\sharp \circ \star = \star \circ \sharp$ is equivalent to (3.15), square relation (3.13) and use $\mathcal{D}^2 = \overline{\mathcal{D}}^2 = \pm \mathbf{1}$ to deduce that $\text{const} = \pm 1$. We cannot have $\text{const} = -1$ because $\overline{\mathcal{D}} = -C^t \mathcal{D} C^t$ and $\mathcal{D}^t C \mathcal{D} = C$ would imply $\overline{\mathcal{D}} = -\mathcal{D}^{t-1}$ i.e. $\overline{\mathcal{D}} \mathcal{D}^t = -\mathbf{1}$ that is absurd ($\sum_j \overline{\mathcal{D}}_{ij} \mathcal{D}_{ij} \neq -1$). Similarly

$\overline{\mathcal{D}} = \pm \mathcal{D}$ is equivalent to $\sharp \circ \times = \times \circ \sharp$ [just consider $\mathbf{1}$ instead of C^t in (3.13)]. Since $\overline{\mathcal{D}}\mathcal{D} = \mathbf{1}$, $\overline{\mathcal{D}} = \mathcal{D}$ is only compatible with $\mathcal{D}^2 = \mathbf{1}$, while $\overline{\mathcal{D}} = -\mathcal{D}$ is only compatible with $\mathcal{D}^2 = -\mathbf{1}$. Finally the q -plane map defined by $x^\sharp = \mathcal{D}x$ is an involutive automorphism of the quantum plane algebra if and only if $\mathcal{D}^2 = \mathbf{1}$. In this case $\overline{\mathcal{D}} = \mathcal{D}$ (for $\times$) or $C^t\overline{\mathcal{D}} = \mathcal{D}C^t$ (for $\star$) so that $\sharp$ commutes with $\times$ or $\star$ on the q -plane. Compatibility of $\times$ and $\star$ with the quantum group coaction δ : $\delta(x^\times) = T^\times \otimes x^\times$ and $\delta(x^\star) = T^\star \otimes x^\star$ is straightforward. $\square\square\square$

Note 1 We can interpret the $\mathcal{D}$ matrix of Theorem 1 as a special T matrix with complex entries. Then conditions (3.11), (3.12) and $\mathcal{D}^2 = \pm \mathbf{1}$ simply state that $\mathcal{D}$ is a quantum orthogonal matrix, moreover the constraints (3.15) respectively state that $\mathcal{D}$ satisfies (modulo a minus sign in the $\times$ case) the reality conditions $T^\times = T$, of $SO_q(n, n)$, $SO_q(n+1, n)$, and $T^\star = C^tTC^t$ of $SO_q(N, \mathbf{R})$.

In order to study the real forms related to $\times^\sharp$ and $\star^\sharp$, where $\sharp$ is given by (3.9), (3.10), we find a linear transformation $x \rightarrow x' = Mx$, $T \rightarrow T' = MTM^{-1}$ such that the new coordinates x' and T' are real and the new metric $C' = (M^{-1})^tCM^{-1}$, at least in the $q \rightarrow 1$ limit, becomes diagonal.² Then the signature of the metric identifies the real form associated to the given $\star$ -conjugation.

The transformation matrices M and the corresponding transformed metrics in the $q = 1$ limit for $\star$ and $\star^\sharp$ explicitly read:

$$\begin{array}{cc}
N \text{ even} & , \quad N \text{ odd} \\
\left. \begin{array}{l}
M = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & & & & & & & 1 \\ & \dots & & & & & & \\ & & 1 & & & & & 1 \\ & & & 1 & 1 & & & \\ & & & i & -i & & & \\ & & & & & i & -i & \\ & & & & & & \dots & \\ i & & & & & & & -i \end{pmatrix} \\
C' = \text{diag}(1, \dots, 1, 1, 1, 1, \dots, 1)
\end{array} \right\} \text{for } \star
\end{array}
\quad , \quad
\left. \begin{array}{l}
M = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & & & & & & & 1 \\ & \dots & & & & & & \\ & & 1 & & & & & 1 \\ & & & \sqrt{2} & & & & \\ & & & i & -i & & & \\ & & & & & \dots & & \\ i & & & & & & & -i \end{pmatrix} \\
C' = \text{diag}(1, \dots, 1, 1, 1, 1, \dots, 1)
\end{array} \right\} \text{for } \star
\end{array}$$

$$\begin{array}{cc}
\left. \begin{array}{l}
M = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & & & & & & & 1 \\ & \dots & & & & & & \\ & & 1 & & & & & 1 \\ & & & 1 & 1 & & & \\ & & & -1 & 1 & & & \\ & & & i & & -1 & & \\ & & & & & & \dots & \\ i & & & & & & & -i \end{pmatrix} \\
C' = \text{diag}(1, \dots, 1, 1, -1, 1, \dots, 1)
\end{array} \right\} \text{for } \star^\sharp
\end{array}
\quad , \quad
\left. \begin{array}{l}
M = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & & & & & & & 1 \\ & \dots & & & & & & \\ & & 1 & & & & & 1 \\ & & & i\sqrt{2} & & & & \\ & & & i & -i & & & \\ & & & & & \dots & & \\ i & & & & & & & -i \end{pmatrix} \\
C' = \text{diag}(1, \dots, 1, -1, 1, \dots, 1)
\end{array} \right\} \text{for } \star^\sharp
\end{array}$$

²For $q \neq 1$ it is not possible to have a diagonal metric, indeed it is not possible to diagonalize C via the transformation $C' = (M^{-1})^tCM^{-1}$ because $C^t \neq C \Rightarrow C'^t \neq C'$. Of course in the $q \rightarrow 1$ limit $C^t = C$.

Therefore $\star$ gives the real form $SO_q(N, \mathbf{R})$, while $\star^\sharp$ gives the real form $SO_q(N - 1, 1)$. The matrices corresponding to the $\times^\sharp$ conjugation have a similar structure with nonzero entries only in the diagonals. For N even we obtain the real form $SO_q(n + 1, n - 1)$ with $|q| = 1$. For N odd we obtain the real form $SO_q(n, n + 1)$ with $|q| = 1$; this is isomorphic to the real form $SO_q(n + 1, n)$.

Proof: We construct an automorphism α of $SO_q(N)$ such that $\alpha \circ \times^\sharp = \times \circ \alpha$. Consider

$$\alpha : SO_q(N) \rightarrow SO_q(N), \quad \alpha(T) \equiv ATA^{-1}, \quad A \equiv \text{diag}(i, \dots, i, 1, -i, \dots, -i) \quad (3.17)$$

The matrix A (like $\mathcal{D}$) satisfies (3.11) and (3.12) so that $R\alpha(T_1)\alpha(T_2) = \alpha(T_2)\alpha(T_1)R$, $\alpha(T)^t C \alpha(T) = C$; therefore α is an automorphism of $SO_q(N)$. We also have $\mathcal{D}A = -\overline{A}$ so that $\alpha(T^{\times^\sharp}) = [\alpha(T)]^\times$ indeed:

$$\alpha(T^{\times^\sharp}) = \mathcal{D}\alpha(T)\mathcal{D}^{-1} = \mathcal{D}ATA^{-1}\mathcal{D}^{-1} = \overline{A}T\overline{A}^{-1} = (ATA^{-1})^\times = [\alpha(T)]^\times. \quad (3.18) \quad \square$$

We now find real forms $SO_q(l, m)$ with l and m arbitrary by considering other examples of $\mathcal{D}$ matrices that, as in Theorem 1, satisfy (3.11), (3.12), $\mathcal{D}^2 = \mathbf{1}$ and $\overline{\mathcal{D}} = \mathcal{D}$ (for $\times$) or $\overline{\mathcal{D}} = C^t \mathcal{D} C^t$ (for $\star$). Following [2] we first study the matrices $\mathcal{D}' = \text{diag}(\varepsilon_1, \dots, \varepsilon_N)$, where $\varepsilon_j^2 = 1$, $\varepsilon_{j'} = \varepsilon_j$, $j = 1, \dots, N$ and $\varepsilon_{n+1} = 1$ for $N = 2n + 1$ or $\varepsilon_n = \varepsilon_{n+1} = 1$ for $N = 2n$. [In the even case $\varepsilon_j = -1 \forall j < n$ gives $-\text{diag}(1, \dots, 1, -\varepsilon_n, -\varepsilon_{n+1}, 1, \dots, 1)$ that differs from minus the identity only if $\varepsilon_n = \varepsilon_{n+1} = 1$, this is why we impose $\varepsilon_n = \varepsilon_{n+1} = 1$. Similarly in the odd case, where moreover $\varepsilon_j = -1 \forall j < n + 1$ gives minus the $\mathcal{D}$ matrix in (3.10)]. The involutive automorphisms $\natural$ associated to the $\mathcal{D}'$ matrices are given by $T^\natural = \mathcal{D}' T \mathcal{D}'^{-1}$; the corresponding conjugations are $\times^\natural$ and $\star^\natural$. Since the $\mathcal{D}'$ matrices commute also with the $\mathcal{D}$ matrices in (3.10) we have $(\mathcal{D}\mathcal{D}')^2 = \mathbf{1}$, $\sharp^\natural$ is an involutive automorphism, $\times \circ \sharp^\natural = \sharp^\natural \circ \times$, $\star \circ \sharp^\natural = \sharp^\natural \circ \star$ and the compositions $\times^\natural \circ \sharp^\natural$, $\star^\natural \circ \sharp^\natural$ are again conjugations. This holds both in the quantum group and in quantum plane case.

The conjugations $\times^\natural$ and $\star^\natural$ still give the real forms $SO_q(n + 1, n)$, $SO_q(n, n)$, $SO_q(n + 1, n - 1)$ with $|q| = 1$.

Proof: rewrite (3.18) for the automorphism α given by $A = \text{diag}(\sigma(\varepsilon_1), \dots, \sigma(\varepsilon_N))$ where $\sigma(\varepsilon_j) = 1$ if $\varepsilon_j = 1$, $\sigma(\varepsilon_j) = i$ if $\varepsilon_j = -1$ and $j < j'$, $\sigma(\varepsilon_j) = -i$ if $\varepsilon_j = -1$ and $j > j'$. $\square$

The conjugations $\star^\natural$, studied in [2], give the real forms $SO_q(l, m)$ where, if $l + m$ is odd, l and m are arbitrary numbers satisfying the only constraint $l + m = 2n + 1$, if $l + m$ is even l and m are both even. Finally real forms $SO_q(l, m)$ with $l + m = 2n$ and l and m both odd are given by the conjugations $\star^\sharp$.

In the $SO_q(2n + 1)$ case there are 2^n matrices $\mathcal{D}'$ and we conclude that the number of real forms $SO_q(2n + 1, \mathcal{D}')$ of $SO_q(2n + 1)$ is 2^n for $q \in \mathbf{R}$. For $|q| = 1$ we have just the real form $SO_q(n + 1, n)$. In [5] Twietmeyer classifies real forms of $\mathcal{U}_q(g)$ where g is semisimple (and more in general symmetrizable Kac-Moody); there are 2^n real forms of $\mathcal{U}_q(so(2n + 1))$ for $q \in \mathbf{R}$ and just one real form for $|q| = 1$. Comparison with [5] shows that via $\times$, $\star$ and the $\mathcal{D}'$ matrices we have described all real forms of $SO_q(2n + 1)$.

One proceed similarly in the $SO_q(2n)$ case. We have 2^{n-1} matrices $\mathcal{D}'$ that give 2^{n-1} $\star^{\natural}$ -conjugations. The conjugations $\star^{\natural}$ are only 2^{n-2} since the matrices $\mathcal{D}'_1 = \text{diag}(\varepsilon_1, \dots, \varepsilon_{n-1}, 1, 1, \varepsilon_{n+2}, \dots, \varepsilon_{2n})$ and $\mathcal{D}'_2 = \text{diag}(-\varepsilon_1, \dots, -\varepsilon_{n-1}, 1, 1, -\varepsilon_{n+2}, \dots, -\varepsilon_{2n})$ give equivalent conjugations $\star^{\natural_1}$ and $\star^{\natural_2}$.

Proof: The map $\alpha(T) = ATA^{-1}$ given by $A = \text{diag}(-1, \dots, -1, 1, \dots, 1)$ is an automorphism of $SO_q(2n)$. We also have $C^t \mathcal{D} \mathcal{D}'_1 A = \overline{A} C^t \mathcal{D} \mathcal{D}'_2$, where $\mathcal{D}$ is given in (3.10); this implies $\alpha(T^{\star^{\natural_1}}) = [\alpha(T)]^{\star^{\natural_2}}$:

$$\begin{aligned} \alpha(T^{\star^{\natural_1}}) &= \alpha(C^t \mathcal{D} \mathcal{D}'_1 T \mathcal{D}'_1 \mathcal{D} C^t) = (C^t \mathcal{D} \mathcal{D}'_1 A) T (A^{-1} \mathcal{D}'_1 \mathcal{D} C^t) \\ [\alpha(T)]^{\star^{\natural_2}} &= \overline{A} T^{\star^{\natural_2}} \overline{A}^{-1} = (\overline{A} C^t \mathcal{D} \mathcal{D}'_2) T (\mathcal{D}'_2 \mathcal{D} C^t \overline{A}^{-1}) . \end{aligned} \quad \square$$

We now study quantum deformations of $SO^*(2n)$. The real form $SO^*(2n)$ in the commutative case is generated by the matrices O that satisfy

$$O^t O = \mathbf{1} \quad , \quad \overline{O}^t J O = J, \quad \text{where } J = \begin{pmatrix} 0 & \mathbf{1}_n \\ -\mathbf{1}_n & 0 \end{pmatrix} . \quad (3.19)$$

Relation $\overline{O}^t J O = J$ is equivalent to $\overline{O} = J O J^{-1}$. There are many real forms $SO_q^*(2n)$, they are described by the matrices $\mathcal{D}'' = i \text{diag}(\varepsilon_1, \dots, \varepsilon_{2n})$ with $\varepsilon_j^2 = 1$, $\varepsilon_{j'} = -\varepsilon_j$, $j = 1, \dots, 2n$ and $\varepsilon_n = -\varepsilon_{n+1} = 1$. It is easy to verify that the matrices $\mathcal{D}''$ satisfy (3.11), (3.12), $\mathcal{D}''^2 = -\mathbf{1}$, (3.15) and that therefore define involutive Hopf algebra automorphisms $\natural : T \mapsto T^{\natural} \equiv \mathcal{D}'' T \mathcal{D}''^{-1}$ that commute with $\star$ and $\times$. The conjugations $\times^{\natural}$ are equivalent to $\times$ [hint: use $A = \text{diag}(e^{-i\varepsilon_1 \pi/4}, \dots, e^{-i\varepsilon_{2n} \pi/4})$] and still give the real form $SO_q(n, n)$ with $|q| = 1$; since $\natural \circ \natural = \text{id}$, the star structures $\times^{\natural}$ are equivalent to $\times$ and still give the real form $SO_q(n+1, n-1)$ with $|q| = 1$.

The conjugations $\star^{\natural}$ give the real forms $SO_q(2n, \mathcal{D}'')$. To prove that all the q -groups $SO_q(2n, \mathcal{D}'')$ are deformations of $SO^*(2n)$ we show that in the commutative limit all the conjugations $\star^{\natural}$ are equivalent. Then we find the transformation to the basis (3.19).

Proof: $\star^{\natural}$ given by $\mathcal{D}'' = i \text{diag}(\varepsilon_1, \dots, \varepsilon_{2n})$, for $q = 1$, is equivalent to the conjugation given by $\mathcal{D}'_1 = i \text{diag}(1, \dots, 1, -1, \dots, -1)$ via the automorphism $\alpha(T) = ATA^{-1}$ where A is defined in (3.20). Notice that only in the $q = 1$ limit A satisfies (3.11) and (3.12).

$$A = \frac{1}{2} \begin{pmatrix} 1+\varepsilon_1 & & & & & & 1-\varepsilon_1 \\ & \dots & & & & & \\ & & 1+\varepsilon_{n-1} & & 1-\varepsilon_{n-1} & & \\ & & & 2 & 0 & & \\ & & & 0 & 2 & & \\ & & & & & 1-\varepsilon_{n-1} & 1+\varepsilon_{n-1} \\ & & & & & & \dots \\ 1-\varepsilon_1 & & & & & & 1+\varepsilon_1 \end{pmatrix}, \quad M'' = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & & & & & & 1 \\ & \dots & & & & & \\ & & 1 & & & 1 & \\ & & & 1 & 1 & & \\ i & & & & & & -i \\ & i & & & & & -i \\ & & \dots & & \dots & & \\ & & & i & -i & & \end{pmatrix} \quad (3.20)$$

To obtain (3.19) consider $\mathcal{D}'_1 = i \text{diag}(1, \dots, 1, -1, \dots, -1)$, in the classical limit and in the basis $O = M'' T M''^{-1}$, where M'' is defined in (3.20), relation $T^t C T = C$ reads $O^t O = \mathbf{1}$ and relation $T^{\star^{\natural}} = C^t \mathcal{D}'' T \mathcal{D}''^{-1} C^t$ reads $\overline{O} = J O J^{-1}$. [Hint: $\overline{O} = O^{\star^{\natural}} = \overline{M''} T^{\star^{\natural}} \overline{M''}^{-1}$]. $\square$

In conclusion in the $SO_q(2n)$ case, for $q \in \mathbf{R}$, there are $2^{n-1} + 2^{n-2}$ real forms $SO_q(2n, \mathcal{D}')$ and 2^{n-1} real forms $SO_q(2n, \mathcal{D}'')$ for a total of $2^n + 2^{n-2}$ real forms as in [5]. Also, as in [5], for $|q| = 1$ we have only 2 inequivalent conjugations.

We have thus constructed all real forms of $SO_q(N)$.

Note 2 In the classical case real forms of $so(N)$ (and more in general of a semisimple Lie algebra) are classified via involutive automorphisms of the compact real form $so(N, \mathbf{R})$. (See for example [9]). The procedure we have followed to describe the real forms of $SO_q(N)$ is exactly the same as that of the classical case reformulated in the $*$ -structure language. More explicitly, we recall that an involutive automorphism $\sharp$ of the compact form g_{ct} of the semisimple complex Lie algebra $g = g_{ct} \oplus i g_{ct}$, splits g_{ct} in the direct sum $g_{ct} = t \oplus p$ where t, p are respectively the $\sharp$ eigenspaces with eigenvalue $+1, -1$. Then $g_{\mathbf{R}} = t \oplus i p$, is a non-compact real form of g . If $\star$ is the star structure associated to g_{ct} : $g_{ct} = \{\chi \in g \mid \chi^\star = -\chi\}$ (recall Section 2), then we have $g_{\mathbf{R}} = t \oplus i p = \{\chi \in g \mid \chi^{\star\sharp} = -\chi\}$; this last relation shows that $\star^\sharp$ is the star structure $*$ canonically associated to the real form $g_{\mathbf{R}}$. It naturally extends to $\mathcal{U}(g)$ so that

$$* = \sharp \circ \star \quad \text{on } \mathcal{U}(g) . \quad (3.21)$$

Since all real forms can be found via involutive automorphisms of g_{ct} , all $*$ -structures can be found via (3.21) i.e. via composition of the compact star structure $\star$ with involutive automorphisms of g_{ct} . Notice that $\sharp$ is an involutive automorphism of g_{ct} if and only if $\sharp$ is an involutive automorphism of $g = g_{ct} \oplus i g_{ct}$ and $\sharp \circ \star = \star \circ \sharp$.

In the orthogonal case (for $N \neq 8$) all automorphisms are realized, in the defining representation of $g_{ct} = so(N, \mathbf{R})$, via an orthogonal matrix $\mathcal{D} \in O(N, \mathbf{R})$ (with $\mathcal{D}^2 = \pm \mathbf{1}$ if the automorphism is involutive). Explicitly, let $\{\chi_i\}$ be a basis of g_{ct} , we have $\sharp: \chi_i \mapsto \chi_i^\sharp \equiv \mathcal{D} \chi_i \mathcal{D}^{-1} \equiv \mathbf{D}_i^j \chi_j$. We can use the duality (2.7) to deduce how $*$ in (3.21) acts on the functionals $T^a_b \in \mathcal{F}(G)$ [$G = SO(N, \mathbf{C})$]. More simply, apply (3.21) to $T^a_b(g) = (e^{x^i(g)\chi_i})^a_b$ where $g \in G$ and $x^i(g)$ are the (complex) coordinates of g ; we obtain

$$T^* = T^{\star\sharp} = \mathcal{D} T^* \mathcal{D}^{-1} . \quad (3.22)$$

This formula shows that in the classical case all $*$ -structures of $\mathcal{F}(G)$ are given via involutive automorphisms of $g_{ct} = so(N, \mathbf{R})$, i.e., via involutive automorphisms of $Fun(SO(N, \mathbf{R}))$; they are realized via matrices $\mathcal{D} \in O(N, \mathbf{R})$. (For $N = 8$ all $*$ -structures are given by (3.22) as well). The same occurs in the quantum case. The $*$ -structures $\star^\sharp$, $\star^\natural$ and $\star^{\sharp\sharp}$ we have considered have all the form (3.22) and correspond to the involutive automorphisms $\sharp$, $\natural$ and $\sharp\sharp$ of $SO_q(N)$ that commute with $\star$ (see Theorem 1), i.e. correspond to the involutive automorphisms of the compact form $SO_q(N, \mathbf{R})$. $\sharp$, $\natural$ and $\sharp\sharp$ are realized via the matrices $\mathcal{D}', \mathcal{D}'', \mathcal{D}\mathcal{D}'$ [$\mathcal{D}$ here is given in (3.10)]; these are compact quantum orthogonal matrices, i.e. are matrices with complex entries that satisfy the orthogonality and reality relations: (2.1), (2.2), (3.15) [cf. Note 1] and have q -determinant $= \pm 1$.

We now show that $\mathcal{D}', \mathcal{D}'', \mathcal{D}\mathcal{D}'$ classify the involutive automorphisms of $SO_q(N, \mathbf{R})$ ($\forall N \neq 8$). From [5] we deduce that $SO_q(2n+1, \mathbf{R})$ has 2^n involutive automorphisms. The involutive automorphisms given by $T^\natural = \mathcal{D}'T\mathcal{D}'^{-1}$ [$\mathcal{D}' = \text{diag}(\varepsilon_1, \dots, \varepsilon_{2n+1})$; recall that $-\mathcal{D}$, where $\mathcal{D}$ is defined in (3.10) is a particular $\mathcal{D}'$ matrix] are 2^n and we conclude that these are all the involutive automorphisms of $SO_q(2n+1, \mathbf{R})$. In the classical case all automorphisms of $so(2n+1, \mathbf{R})$ are inner i.e. can be realized via matrices of $SO(2n+1, \mathbf{R})$, similarly, in the quantum case, the matrices $\mathcal{D}' = \text{diag}(\varepsilon_1, \dots, \varepsilon_{2n+1})$ satisfy the defining relations of the compact quantum group $SO_q(2n+1, \mathbf{R})$.

The automorphism group of $so(2n, \mathbf{R})$ (for $n \neq 4$) has two disconnected components and is $O(2n, \mathbf{R})$. This corresponds to the discrete $\mathbf{Z}_2$ symmetry of the Dynkin diagram of the D_n series. In the quantum case there are 2^n [5] involutive inner automorphisms. These are all realized via the 2^{n-1} matrices $\mathcal{D}'$ and the 2^{n-1} matrices $\mathcal{D}''$; these matrices satisfy the defining relations of $SO_q(N, \mathbf{R})$. On the other hand there are 2^{n-1} outer involutive automorphisms, these are all realized via the 2^{n-1} matrices $\mathcal{D}\mathcal{D}'$ that are quantum orthogonal matrices with $\det_q \mathcal{D}\mathcal{D}' = -1$.

Note 3 There are two different Lorentz groups obtained as real forms of $SO_q(4)$: $SO_q(3, 1)$ with $q \in \mathbf{R}$ and $SO_q(3, 1)$ with $|q| = 1$. More in general we have deformations of $SO(n+1, n)$, $SO(n, n)$, $SO(n+1, n-1)$ both for $|q| = 1$ and for $q \in \mathbf{R}$.

Note 4 As shown in [8] the classical embedding of $SO(N)$ in $SO(N+2)$ holds also in the quantum case. We have that the universal enveloping algebra $\mathcal{U}_q(so(N))$ of $SO_q(N)$ is a Hopf subalgebra of $\mathcal{U}_q(so(N+2))$. Otherwise stated $SO_q(N)$, that is dual to $\mathcal{U}_q(so(N))$, is a quotient of $SO(N+2)$ via a Hopf ideal H : $SO_q(N) = SO(N+2)/H$. In the commutative case H is the ideal in $Fun(SO(N+2))$ generated by the equivalence relation $f \sim f' \Leftrightarrow f|_{SO(N)} = f'|_{SO(N)}$ where f and f' are generic functions on $SO(N+2)$ and $f|_{SO(N)}$ is f restricted to $SO(N)$.

On the other hand the classical embedding of $SO(N)$ in $SO(N+1)$ does not occur at the quantum level. In particular the q -Lorentz groups discussed here do not “contain” three dimensional q -orthogonal subgroups, i.e., $SO_q(3)$ cannot be found as a quotient $SO_q(4)/H$ with respect to an appropriate ideal H .

It is interesting to note that however the q -Minkowski plane for $q \in \mathbf{R}$ “contains” the euclidean q -plane and the three dimensional q -Minkowski plane.

Proof: The q -Minkowski coordinates satisfy:

$$\begin{aligned} x^1 x^2 &= q x^2 x^1, \quad x^1 x^3 = q x^3 x^1, \quad x^2 x^4 = q x^4 x^2, \quad x^3 x^4 = q x^4 x^3, \\ x^2 x^3 &= x^3 x^2, \quad x^1 x^4 = x^4 x^1 - (q - q^{-1}) x^2 x^3. \end{aligned} \quad (3.23)$$

with $(x^1)^{\star\sharp} = q x^4$, $(x^2)^{\star\sharp} = x^2$ and $(x^3)^{\star\sharp} = x^3$. The $SO_q(3)$ plane commutations are

$$y^1 y^2 = q y^2 y^1, \quad y^2 y^3 = q y^3 y^2, \quad y^1 y^3 = y^3 y^1 - (q^{\frac{1}{2}} - q^{-\frac{1}{2}}) y^2 y^2 \quad (3.24)$$

The conjugation that gives the euclidean plane is $(y^1)^\star = q y^3$, $(y^2)^\star = y^2$, while for the $SO_q(2, 1)$ plane we have $(y^1)^{\star\sharp} = q y^3$, $(y^2)^{\star\sharp} = -y^2$. To obtain the $SO_q(3, \mathbf{R})$

plane from (3.23) we quotient with respect to the relation $x^2 = x^3$ and identify $x^1, \sqrt{q^{\frac{1}{2}} + q^{-\frac{1}{2}}}x^2, x^4$ respectively with y^1, y^2, y^3 . To obtain the $SO_q(2, 1)$ plane we impose the relation $x^2 = -x^3$ and identify $x^1, i\sqrt{q^{\frac{1}{2}} + q^{-\frac{1}{2}}}x^2, x^4$ respectively with y^1, y^2, y^3 . $\square$

One similarly proves that the $|q| = 1$ $SO_q(2, 1)$ plane is obtained from the $|q| = 1$ q -Minkowski spacetime with $\times^\sharp$ -conjugation (3.16): $(x^1)^{\times^\sharp} = x^1, (x^2)^{\times^\sharp} = x^3, (x^4)^{\times^\sharp} = x^4$ (cf. [3]), via the quotient $x^2 = x^3$. Note that this $|q| = 1$ q -Minkowski spacetime has also been studied in [7], there the q -commutations and the conjugation are derived from the conformal group $SU_q(2, 2)$. Explicitly, in [7] the conjugation used is $\times^{\sharp\sharp}$ with $\mathcal{D}' = \text{diag}(-1, 1, 1, -1)$: $(x^1)^{\times^{\sharp\sharp}} = -x^1, (x^2)^{\times^{\sharp\sharp}} = x^3, (x^4)^{\times^{\sharp\sharp}} = -x^4$, this conjugation we have proved to be equivalent to $\times^\sharp$.

Acknowledgments

The autor thanks Leonardo Castellani for fruitful discussions and encouragement. The autor thanks Bogdan Morariu, Nicolai Reshetikhin and Bruno Zumino for their valuable comments and suggestions.

This work has been supported by an INFN post-doctoral grant (concorso n. 6077/96). It has been accomplished through the Director, Office of Energy Research, Office of High Energy and Nuclear Physics, Division of High Energy Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098 and by the National Science Foundation under grant PHY-95-14797.

References

- [1] B. L. Cerchiai, J. Wess, *q-Deformed Minkowski Space based on a q-Lorentz Algebra* math.QA/9801104; J. Wess, *Quantum groups and q-lattices in phase-space*, in the Proceedings of the 5-th Hellenic School and Workshop on Elementary Particle Physics, Corfu, September 1995, q-alg/9607002.
- [2] L.D. Faddeev, N.Yu. Reshetikhin and L.A. Takhtajan, *Quantization of Lie Groups and Lie Algebras*, Algebra i Anal. **1** **1** (1989) 178 (Leningrad Math. J. **1** 193 (1990)).
- [3] E. Celeghini, R. Giachetti, A. Reyman, E. Sorace, M. Tarlini, *$SO_q(n+1, N-1)$ as a Real Form of $SO_q(2n, \mathbb{C})$* , Lett. Math. Phys. **23** (1991) 45.
- [4] V.K. Dobrev, *Canonical q-Deformations of Noncompact Lie (Super-) Algebras*, J. Phys. A: Math. Gen. **26** (1993) 1317, (Göttingen University preprint, July 1991).
- [5] E. Twietmeyer, *Real Forms of $\mathcal{U}_q(g)$* , Lett. Math. Phys. **24** (1992) 49.

- [6] G. Fiore, J. Phys. *Quantum Groups* $SO_q(N)$, $Sp_q(N)$ have determinants, too **A27** (1994) 3795.
- [7] V.K. Dobrev, *New q - Minkowski space-time and q - Maxwell equations hierarchy from q -conformal invariance* Phys. Lett. **341B** 133 (1994), erratum -ibid. **346B** 427 (1995).
- [8] P. Aschieri and L. Castellani, *Universal enveloping algebra and differential calculi on inhomogeneous orthogonal q -groups*, LBNL 40330, q-alg/9705023. to be publ. in J. Geom. and Phys.
- [9] R. Gilmore *Lie Groups, Lie algebras and Some of Their Applications* Wiley-Interscience New York - London - Sydney - Toronto (1974). M. Hausner, J. Schwartz, *Lie Groups; Lie Algebras* Gordon and Breach, New York - London - Paris (1968).